

Case Number: 167968

Gender: M

Data Entry Date: 30/08/2001

Weight (kg):

Hospitalisation:

Age: 13Y

Onset Date: 10/07/2001

DOB: 05/04/1988

Outcome:

Causality: Causality possible

Unknown

Information: Following ingestion of methylphenidate attenta brand only (ritalin - no adverse effects).

Reaction:

Preferred Term

Severity

Report Description

Treatment

Pain Migraine Severe Severe pain.

With holding attenta when

possible.

Vomiting

Medicine details:

ATTENTA (Suspected)

Reason:

30 Milligram

Daily

Batch:

Started: 10/07/2001

Stopped:

PUSIME02 Page 1 of 5



Case Number: 169403

Gender: M

Data Entry Date: 22/10/2001 **Weight (kg)**: 16

Hospitalisation: Age: 06Y

Onset Date: 10/08/2001 **DOB**: 08/09/1994

Outcome: Causality: Causality probable

Recovered

Information:

Reaction:

Preferred Term Severity Report Description Treatment

Aggression Agitation

Depression Required a visit to doctor, date

of recovery 4.9.01. immediate change once was back on

ritalin.

Medicine details:

ATTENTA (Suspected) Reason: Behavior disorders of childhod

20 Milligram Daily

Batch: **Started**: 10/08/2001 **Stopped**: 04/09/2001

PUSIME02 Page 2 of 5



Case Number: 181914

 $\textbf{Gender}: \ M$

Data Entry Date: 22/01/2003

Weight (kg): 0

Hospitalisation:

Age: 16Y

Onset Date:

DOB:

Outcome:

Causality: Causality possible

Recovered

Information:

Reaction:

Preferred Term

Severity Report Description

Treatment

Drug ineffective

Decreased control of

Changed back to the Ritalin

behaviour.

Brand.

Abnormal behaviour

Medicine details:

ATTENTA (Suspected)

Reason : Behavior disorders of childhod

Tablet

40 Milligram

Daily

Oral

Batch:

Started:

Stopped:

PUSIME02 Page 3 of 5



Case Number: 232021

Gender: M

Data Entry Date: 15/08/2007

Weight (kg): 0

Hospitalisation: Required a visit to the doctor

Age:

Onset Date: 15/04/2007

DOB: 02/06/1967

Outcome:

Causality: Causality certain

Not yet recovered

Information: CERT(RECHALL)

Reaction :			
Preferred Term	Severity	Report Description	Treatment
Gastrointestinal inflammation	Caused or prolonged inpatient hospitalisation	Patient developed bilateral upper quadrant abdominal pain. on 2/5/07 developed symptoms of intestinal obstruction and underwent laparotomy and found his bowel was inflamed. Attenta was avoided until mid July 2007 when restarted medication. Within one day the patient developed severe abdominal pain, angieodema of his lips and face, sweating, poly dipsia and polyuria. Attenta was discontinued again until 25/7/07 when patient took a tablet 10mg twice daily the day before his appointment to demonstrate the perceived side effects of his treatment with Attenta. The adverse reactions occured.	laparotomy, ceased Attenta.
Chest discomfort		Tightness in chest, tightness in the stomach area, lumps in the arms below the elbow, muscle pain, hot sweats, nausea. Drug ineffective.	Patient decreased dose of Ritalin himself and then stopped treatment.
Abdominal pain	Caused or prolonged inpatient hospitalisation		
Angioedema	Caused or prolonged inpatient hospitalisation		
Blood pressure increased	Caused or prolonged inpatient hospitalisation		
Hyperhidrosis	Caused or prolonged inpatient hospitalisation		

PUSIME02 Page 4 of 5 Reaction:

Preferred Term Severity Report Description Treatment

Polydipsia Caused or prolonged

inpatient

hospitalisation

Polyuria Caused or prolonged

inpatient

hospitalisation

Tachycardia Caused or prolonged

inpatient hospitalisation

Medicine details:

ATTENTA (Suspected) Reason :

Batch: Started: Stopped: 0

Laboratory Investigations:

Type Range Date Tested Result Details

Blood pressure 170/100 and pulse of 110 beats per

minute.

Selection Parameters: Date Range: 01/01/1960 To 31/12/2059 Unclear causality excluded GM medicines Only Medicine Names: ATTENTA

PUSIME02 Page 5 of 5